arXiv:1603.01502v1 [math.PR] 4 Mar 2016

Multidimensional Lévy White Noises in Weighted Besov Spaces

March 7, 2016

Julien Fageot, Alireza Fallah, and Michael Unser

Abstract

In this paper, we study the Besov regularity of d-dimensional Lévy white noises. More precisely, we describe new sample paths properties of a given white noise in terms of weighted Besov spaces. In particular, the smoothness and integrability properties of Lévy white noises are characterized using the Blumenthal-Getoor indices. Our techniques rely on wavelet methods and generalized moments estimates for Lévy white noises.

1 Introduction

This paper is dedicated to the study of the regularity of a general d-dimensional Lévy white noise in terms of Besov spaces and is a continuation of our previous work [13]. A random process is traditionally defined as a collection (X_t) of random variables indexed by $t \in \mathbb{R}$, with some adequate properties. For instance, Lévy processes are described as stochastically continuous random processes with independent and stationary increments [1, 35]. However, it is not possible to define Lévy white noises in the traditional framework. Typically, in the 1D setting, it is tempting to introduce Lévy white noises as the derivatives of Lévy processes. The well-known issue is that Lévy processes are (almost surely) not differentiable in the classical sense: The derivative of a non-trivial Lévy process does not have a pointwise interpretation. An alternative way of introducing random processes is based on the abstract theory of measures on function spaces, as developed by Bogachev [5] among others. In this context, a random process is a random variable that takes values in a function space endowed with the adequate measurable structure. In this spirit, Gelfand [15] and Itô [21] have independently introduced the concept of generalized random processes, defined as random elements in the Schwartz space of generalized functions [40]. This approach was more extensively exposed in [16, Chapter 3] and [22]. The d-dimensional Schwartz space has the advantage of being stable by (weak) differentiation, and does not only include not only the d-dimensional Lévy white noises, but also all their derivatives.

Measuring regularity with Besov spaces Since we are considering processes that have no pointwise interpretation, we should consider function spaces with negative smoothness. When talking about the regularity of random processes, the Sobolev or the Hölder regularities are natural concepts that comes into mind. In order to be more general, we will question the Besov regularity of Lévy white noises. Besov spaces include both Sobolev and Hölder spaces, and provide a finer measure of the regularity of a function [44, 45]. Evaluating the Besov regularity of random processes over $\mathbb{R}^d$ requires the introduction of weights, since they are generally not decreasing towards infinity. Thereafter, we therefore consider weighted Besov spaces or local Besov spaces.

Regularity of Lévy white noises and related processes To the best of our knowledge, the Besov regularity of d-dimensional Lévy white noises has never been addressed in all generality. Kusuoka [25] estimated the weighted Sobolev regularity of the Gaussian white noise, while Veraar [48] obtained complete results on the local Besov regularity of the Gaussian white noise. However, these works are based on intrinsic Gaussian methods and are not easily extended to the non-Gaussian case. In [13], we derived new results on the Besov regularity of symmetric-α-stable (SαS) Lévy white noises on the d-dimensional torus. This paper is an extension of [13] in two ways: (1) we consider Lévy white noises over $\mathbb{R}^d$ and deduce the local results as corollaries, and (2) we extend the results from SαS white noises to general Lévy white noises.

Other important works on the Besov regularity of 1-dimensional Lévy processes shall be mentioned. The pioneer works concern the Brownian motion [3, 8, 33]; see also [30] for extensions to more general Gaussian processes, including the fractional Brownian motion. Stable Lévy processes are being studied in [8] and [32]. Note that Rosenbaum [32] is using wavelet techniques similar to ours. The case of general Lévy processes was extensively studied by Schilling, both in the local [36] and weighted cases [37, 38]. Herren obtained similar local results in [18]. In particular, Schilling and Herren rely on the Blumenthal-Getoor indices introduced in [4] for their generalization to non-stable processes. In addition, they extend their results to more general classes of Markov processes. Those indices also play a crucial role in the present study. For a comprehensive survey on the Besov regularity of Lévy processes, we refer the reader to [6].

Contributions and outline The paper is organized as follows.

- In Section 2, we define Lévy white noises in the framework of generalized random processes. We also introduce weighted Besov spaces, based on the wavelet methods developed in [44].
- We derive estimates of the moments of Lévy white noises in Section 3. Moment estimates are standard technical results to study the regularity of Lévy processes [10]. Our contribution is to adapt this question to Lévy white noises by considering general test functions.
- In Section 4, we show that weighted Besov spaces are measurable for the cylindrical σ-field on the space $\mathcal{S}'(\mathbb{R}^d)$ of tempered generalized functions. More concretely, it allows us to address the question of the Besov regularity of any generalized random process.
- Sections 5 and 6 are dedicated to the weighted Besov regularity of d-dimensional Lévy white noises. Section 5 is restricted to weighted L_2-Sobolev spaces, and is therefore an intermediate step required to apply our wavelet techniques in Section 6. The main result of this paper is Theorem 3. There, we give sufficient conditions for a Lévy white noise to be almost surely in a given weighted Besov space in terms of the Blumenthal-Getoor indices of the noise.
- In Section 7, we deduce the local Besov regularity of Lévy white noises from the above.
- Finally, we discuss our results in Section 8 and apply them to some specific families of Lévy white noises that are often encountered in practice.

2 Preliminaries

2.1 Generalized Processes and Lévy White Noises

The stochastic processes of this paper are defined in the framework of generalized random processes [16, Chapter 3]. It allows one in particular to consider Lévy white noises as well-defined

random processes, which is not possible in more traditional approaches since they do not admit a pointwise interpretation.

The Schwartz space of infinitely smooth functions and rapidly decaying functions on $\mathbb{R}^d$ is denoted by $\mathcal{S}(\mathbb{R}^d)$. It is endowed with the topology associated with the following notion of convergence: A sequence (φ_n) of functions in $\mathcal{S}(\mathbb{R}^d)$ converges to $\varphi \in \mathcal{S}(\mathbb{R}^d)$ if, for every multiindex $\boldsymbol{\alpha} \in \mathbb{N}^d$ and every $\mu \geq 0$, the functions $\boldsymbol{x} \mapsto |\boldsymbol{x}|^\mu \mathrm{D}^{\boldsymbol{\alpha}}\{\varphi_n\}(\boldsymbol{x})$ converge to $\boldsymbol{x} \mapsto |\boldsymbol{x}|^\mu \mathrm{D}^{\boldsymbol{\alpha}}\{\varphi\}(\boldsymbol{x})$ in $L_2(\mathbb{R}^d)$, where $|\cdot|$ is the Euclidian norm on $\mathbb{R}^d$. The space $\mathcal{S}(\mathbb{R}^d)$ is a nuclear Fréchet space [42, Section 51]. The topological dual of $\mathcal{S}(\mathbb{R}^d)$ is the space $\mathcal{S}'(\mathbb{R}^d)$ of tempered generalized functions. A *cylindrical set* of $\mathcal{S}'(\mathbb{R}^d)$ is a subset of the form

$$\Big\{u \in \mathcal{S}'(\mathbb{R}^d), \quad (\langle u, \varphi_1\rangle, \ldots, \langle u, \varphi_n\rangle) \in B\Big\}, \tag{1}$$

where $n \geq 1$, $\varphi_1, \ldots, \varphi_n \in \mathcal{S}(\mathbb{R}^d)$, and B is a Borel subset of $\mathbb{R}^n$. We denote by $\mathcal{B}_c(\mathcal{S}'(\mathbb{R}^d))$ the *cylindrical σ-field* of $\mathcal{S}'(\mathbb{R}^d)$, defined as the σ-field generated by the cylindrical sets. Then, $(\mathcal{S}'(\mathbb{R}^d), \mathcal{B}_c(\mathcal{S}'(\mathbb{R}^d)))$ is a measurable space. We fix the probability space $(\Omega, \mathcal{F}, \mathscr{P})$.

Definition 1. A *generalized random process* is a measurable function

$$s \ : \ (\Omega, \mathcal{F}) \to (\mathcal{S}'(\mathbb{R}^d), \mathcal{B}_c(\mathcal{S}'(\mathbb{R}^d))). \tag{2}$$

Its *probability law* is the measure on $\mathcal{S}'(\mathbb{R}^d)$, image of $\mathscr{P}$ by s. For every $B \in \mathcal{B}_c(\mathcal{S}'(\mathbb{R}^d))$,

$$\mathscr{P}_s(B) = \mathscr{P}(\{\omega \in \Omega, s(\omega) \in B\}). \tag{3}$$

The *characteristic functional* of s is defined for every $\varphi \in \mathcal{S}(\mathbb{R}^d)$ by

$$\widehat{\mathscr{P}_s}(\varphi) = \int_{\mathcal{S}'(\mathbb{R}^d)} \mathrm{e}^{\mathrm{i}\langle u, \varphi\rangle} \mathrm{d}\mathscr{P}_s(u). \tag{4}$$

A generalized random process is a random element of the space of tempered generalized functions. The characteristic functional is the infinite-dimensional generalization of the characteristic function. It characterizes the law of s in the sense that

$$\mathscr{P}_{s_1} = \mathscr{P}_{s_2} \Leftrightarrow \widehat{\mathscr{P}}_{s_1} = \widehat{\mathscr{P}}_{s_2}, \tag{5}$$

which we denote by $s_1 \overset{(d)}{=} s_2$ (where (d) stands for equality in distribution). Since the space $\mathcal{S}(\mathbb{R}^d)$ is nuclear, the Minlos-Bochner theorem [16, 29] gives a complete characterization of admissible characteristic functionals.

Theorem 1 (Minlos-Bochner theorem)**.** *A functional $\widehat{\mathscr{P}}$ on $\mathcal{S}(\mathbb{R}^d)$ is the characteristic functional of a generalized random process s if and only if it is continuous and positive-definite over $\mathcal{S}(\mathbb{R}^d)$ and satisfies $\widehat{\mathscr{P}}(0) = 1$.*

Lévy processes are random processes index by $\mathbb{R}$ with stationary and independent increments. They are deeply related to infinitely divisible random variables [35]. For the same reasons, there is a one-to-one correspondence between infinitely divisible laws and Lévy white noises. An infinitely divisible random variable X can be decomposed as $X = X_1 + \cdots + X_n$ for every $n \geq 1$. We say that a function f from $\mathbb{R}$ to $\mathbb{C}$ is a *Lévy exponent* if it is the continuous log-characteristic function of an infinitely divisible random variable. We say moreover that f *satisfies the Schwartz condition* if the moment $\mathbb{E}[|X|^\epsilon]$ of X is finite for some $\epsilon > 0$.

Let $\boldsymbol{X} = (X_1, \ldots, X_n)$ be i.i.d. infinitely divisible random variables with common Lévy exponent f. By independence, the characteristic function of $\boldsymbol{X}$ is

$$\Phi_{\boldsymbol{X}}(\boldsymbol{\xi}) = \exp\left(\sum_{i=1}^{n} f(\xi_i)\right) \tag{6}$$

for every $\boldsymbol{\xi} = (\xi_1, \ldots, \xi_n) \in \mathbb{R}^n$. The class of Lévy white noise can be seen as the generalization of this principle in the continuous domain, up to the replacement of the sum in (6) by an integral.

Definition 2. A *Lévy white noise* is a generalized random process w with characteristic functional of the form

$$\widehat{\mathscr{P}}_w(\varphi) = \exp\left(\int_{\mathbb{R}^d} f(\varphi(\boldsymbol{x}))\mathrm{d}\boldsymbol{x}\right) \tag{7}$$

for every $\varphi \in \mathcal{S}(\mathbb{R}^d)$, where f is a Lévy exponent that satisfies the Schwartz condition.

Gelfand and Vilenkin have proved that the functional (7) is a valid characteristic functional on $\mathcal{D}(\mathbb{R}^d)$, the space of compactly supported and infinitely smooth functions, without the Schwartz condition on f [16]. The Schwartz condition is sufficient to extend this result to $\mathcal{S}(\mathbb{R}^d)$ [12, Theorem 3]. Recently, Dalang and Humeau have shown that this condition is also necessary: A white noise with Lévy exponent that does not satisfy the Schwartz condition is almost surely not in $\mathcal{S}'(\mathbb{R}^d)$ [9, Theorem 3.13].

A Lévy white noise is stationary, in the sense that $w \overset{(d)}{=} w(\cdot - \boldsymbol{x}_0)$ for every $\boldsymbol{x}_0 \in \mathbb{R}^d$. It is moreover independent at every point, meaning that $\langle w, \varphi\rangle$ and $\langle w, \psi\rangle$ are independent whenever φ and $\psi \in \mathcal{S}(\mathbb{R}^d)$ have disjoint supports. In 1-D, we recover the usual notion of white noise, since w is the derivative in the sense of generalized functions of the Lévy process with the same Lévy exponent. This principle can be extended in any dimension $d \geq 2$: The d-dimensional Lévy white noise is the weak derivative $\mathrm{D}_{x_1} \cdots \mathrm{D}_{x_d}\{s\}$ of the d-dimensional Lévy sheet s [9].

2.2 Weighted Sobolev and Besov Spaces

Our goal is to characterize the smoothness of Lévy white noises in terms of weighted Besov spaces. All our results related to Besov spaces would require the corresponding intermediate result for Sobolev spaces which we introduce in Section 2.2.1.

2.2.1 Weighted Sobolev Spaces

We set $\langle \boldsymbol{x}\rangle = \sqrt{1 + |\boldsymbol{x}|^2}$. The Fourier transform of $f \in \mathcal{S}'(\mathbb{R}^d)$ is denoted by $\widehat{f}$. For $\tau \in \mathbb{R}$, we define L_τ (the Bessel operator of order τ) as the pseudo-differential operator with Fourier multiplier $\langle\cdot\rangle^\tau$. In Fourier domain, we write

$$\widehat{\mathrm{L}_\tau\{\varphi\}}(\boldsymbol{\omega}) := \langle\boldsymbol{\omega}\rangle^\tau \widehat{\varphi}(\boldsymbol{\omega}) \tag{8}$$

for every $\boldsymbol{\omega} \in \mathbb{R}^d$ and $\varphi \in \mathcal{S}(\mathbb{R}^d)$. When $\tau > 0$, the operator $\mathrm{I}_\tau = \mathrm{L}_{-\tau}$ is called a Bessel potential [17]. The operator L_τ is self-adjoint, linear, and continuous from $\mathcal{S}(\mathbb{R}^d)$ to $\mathcal{S}(\mathbb{R}^d)$, since its Fourier multiplier is infinitely smooth and bounded by a polynomial function. It can therefore be extended as a linear and continuous operator from $\mathcal{S}'(\mathbb{R}^d)$ to $\mathcal{S}'(\mathbb{R}^d)$.

Definition 3. Let $\tau, \mu \in \mathbb{R}$. The *Sobolev space of smoothness* τ is defined by

$$W_2^\tau(\mathbb{R}^d) := \left\{ f \in \mathcal{S}'(\mathbb{R}^d), \quad \mathrm{L}_\tau\{f\} \in L_2(\mathbb{R}^d) \right\} \tag{9}$$

and the *Sobolev space of smoothness τ and weight μ* is

$$W_2^\tau(\mathbb{R}^d;\mu) := \left\{ f \in \mathcal{S}'(\mathbb{R}^d), \quad \langle\cdot\rangle^\mu f \in W_2^\tau(\mathbb{R}^d) \right\}. \tag{10}$$

We also set $L_2(\mathbb{R}^d;\mu) := W_2^0(\mathbb{R}^d;\mu)$.

We summarize now the basic properties on weighted Sobolev spaces that are useful for our work.

Proposition 1. *The following properties hold for weighted Sobolev spaces.*

- *For $\mu, \tau \in \mathbb{R}$, $W_2^\tau(\mathbb{R}^d;\mu)$ is a Hilbert space for the scalar product*

$$\langle f, g\rangle_{W_2^\tau(\mathbb{R}^d;\mu)} := \langle \mathrm{L}_\tau\{\langle\cdot\rangle^\mu f\}, \mathrm{L}_\tau\{\langle\cdot\rangle^\mu g\}\rangle_{L_2(\mathbb{R}^d)}. \tag{11}$$

  We denote by $\|f\|_{W_2^\tau(\mathbb{R}^d;\mu)} = \langle f, f\rangle^{1/2}_{W_2^\tau(\mathbb{R}^d;\mu)}$ the corresponding norm.

- *For $\mu \in \mathbb{R}$ fixed and for every $\tau_1 \leq \tau_2$, we have the continuous embedding*

$$W_2^{\tau_2}(\mathbb{R}^d;\mu) \subseteq W_2^{\tau_1}(\mathbb{R}^d;\mu). \tag{12}$$

- *For $\tau \in \mathbb{R}$ fixed and for every $\mu_1 \leq \mu_2$, we have the continuous embedding*

$$W_2^\tau(\mathbb{R}^d;\mu_2) \subseteq W_2^\tau(\mathbb{R}^d;\mu_1). \tag{13}$$

- *For $\mu, \tau \in \mathbb{R}$, the operator $\mathrm{L}_{\tau,\mu} : f \mapsto \langle\cdot\rangle^\mu \mathrm{L}_\tau\{f\}$ is an isometry from $L_2(\mathbb{R}^d)$ to $W_2^{-\tau}(\mathbb{R}^d;-\mu)$.*
- *The dual space of $W_2^\tau(\mathbb{R}^d;\mu)$ is $W_2^{-\tau}(\mathbb{R}^d;-\mu)$ for every $\tau, \mu \in \mathbb{R}$.*
- *We have the countable projective limit*

$$\mathcal{S}(\mathbb{R}^d) = \bigcap_{\tau,\mu\in\mathbb{R}} W_2^\tau(\mathbb{R}^d;\mu) = \bigcap_{n\in\mathbb{N}} W_2^n(\mathbb{R}^d;n). \tag{14}$$

- *We have the countable inductive limit*

$$\mathcal{S}'(\mathbb{R}^d) = \bigcup_{\tau,\mu\in\mathbb{R}} W_2^\tau(\mathbb{R}^d;\mu) = \bigcup_{n\in\mathbb{N}} W_2^{-n}(\mathbb{R}^d;-n). \tag{15}$$

Proof. The space $W_2^\tau(\mathbb{R}^d;\mu)$ inherits the Hilbertian structure of $L_2(\mathbb{R}^d)$. For $\tau_1 \leq \tau_2$ and $\mu_1 \leq \mu_2$, we have moreover the inequalities,

$$\|f\|_{W_2^{\tau_1}(\mathbb{R}^d;\mu)} \leq \|f\|_{W_2^{\tau_2}(\mathbb{R}^d;\mu)}, \tag{16}$$

$$\|f\|_{W_2^\tau(\mathbb{R}^d;\mu_1)} \leq \|f\|_{W_2^\tau(\mathbb{R}^d;\mu_2)}, \tag{17}$$

from which we deduce (12) and (13). The relation

$$\|\mathrm{L}_{\tau,\mu} f\|_{W_2^{-\tau}(\mathbb{R}^d;-\mu)} = \|\mathrm{L}_{-\tau}\{\langle\cdot\rangle^{-\mu}\mathrm{L}_{\tau,\mu} f\}\|_{L_2(\mathbb{R}^d)} = \|f\|_{L_2(\mathbb{R}^d)} \tag{18}$$

proves that $\mathrm{L}_{\tau,\mu}$ is an isometry. For every $f, g \in L_2(\mathbb{R}^d)$, we have that

$$\langle \mathrm{L}_\tau\{\langle\cdot\rangle^\mu f\}, \mathrm{L}_{-\tau}\{\langle\cdot\rangle^{-\mu} g\}\rangle_{L_2(\mathbb{R}^d)} = \langle f, g\rangle_{L_2(\mathbb{R}^d)}. \tag{19}$$

Since $W_2^\tau(\mathbb{R}^d;\mu) = \{\mathrm{L}_\tau\{\langle\cdot\rangle^\mu f\},\ f \in L_2(\mathbb{R}^d)\}$, we easily deduce the dual of $W_2^\tau(\mathbb{R}^d;\mu)$ from (19). Finally, we can reformulate the topology on $\mathcal{S}(\mathbb{R}^d)$ as (14). This implies directly (15). □

2.2.2 Weigthed Besov Spaces

Following Triebel [44], our definitions of weighted Besov spaces are based on wavelets. More traditionally, Besov spaces are introduced through the Fourier transform, see for instance [43]. The use of wavelets is equivalent and appears to be more convenient for our framework.

Let us first introduce the wavelet bases that we should use. We denote by $j \geq 0$ the scaling index and $\boldsymbol{m} \in \mathbb{Z}^d$ the shifting index. Consider ψ_{F} and ψ_{M}, which are the father and mother wavelet of a wavelet basis for $L_2(\mathbb{R})$, respectively. We set $\mathrm{G}^0 = \{M, F\}^d$ and $\mathrm{G}^j = \mathrm{G}^0 \backslash (F, \ldots, F)$ for $j \geq 1$. For a gender $G = (G_1, \ldots, G_d) \in \mathrm{G}^0$ and for every $\boldsymbol{x} = (x_1, \ldots, x_d) \in \mathbb{R}^d$, we define

$$\psi_G(\boldsymbol{x}) = \prod_{i=1}^{d} \psi_{G_i}(x_i). \tag{20}$$

Proposition 2 (Section 1.2.1, [44])**.** *For every integer $u \geq 0$, there exist compactly supported wavelets ψ_{F} and ψ_{M} such that*

$$\left\{\psi_{j,G,\boldsymbol{m}}, \quad j \geq 0, G \in \mathrm{G}^j, \boldsymbol{m} \in \mathbb{Z}^d\right\} \tag{21}$$

is an orthonormal basis of $L_2(\mathbb{R}^d)$, where

$$\psi_{j,G,\boldsymbol{m}} := 2^{jd/2} \psi_G(2^j \cdot - \boldsymbol{m}) \tag{22}$$

and ψ_G is defined according to (20)*.*

Concretely, [44] considers separable Daubechies wavelets with the adequate regularity. For $\tau, \mu \in \mathbb{R}$ and $0 < p, q \leq \infty$, the *Besov sequence space* $b^{\tau}_{p,q}(\mu)$ is the collection of sequences

$$\boldsymbol{\lambda} = \{\lambda_{j,G,\boldsymbol{m}}, \quad j \geq 0, G \in \mathrm{G}^j, \boldsymbol{m} \in \mathbb{Z}^d\} \tag{23}$$

such that

$$\|\boldsymbol{\lambda}\|_{b^{\tau}_{p,q}(\mu)} := \left(\sum_{j \geq 0} 2^{jq(\tau - d/p)} \sum_{G \in \mathrm{G}^j} \left(\sum_{\boldsymbol{m} \in \mathbb{Z}^d} \langle 2^{-j} \boldsymbol{m} \rangle^{\mu p} |\lambda_{j,G,\boldsymbol{m}}|^p \right)^{q/p} \right)^{1/q}, \tag{24}$$

with the usual modifications when p and/or $q = \infty$.

Definition 4. Let $\tau, \mu \in \mathbb{R}$ and $0 < p, q \leq \infty$. Fix

$$u > \max(\tau, d(1/p - 1)_+ - \tau) \tag{25}$$

and set $(\psi_{j,G,\boldsymbol{m}})$ a wavelet basis of $L_2(\mathbb{R}^d)$ with regularity u. The *weighted Besov space* $B^{\tau}_{p,q}(\mathbb{R}^d; \mu)$ is the collection of generalized function $f \in \mathcal{S}'(\mathbb{R}^d)$ that can be written as

$$f = \sum_{j,G,\boldsymbol{m}} 2^{-jd/2} \lambda_{j,G,\boldsymbol{m}} \psi_{j,G,\boldsymbol{m}} \tag{26}$$

with $\boldsymbol{\lambda} = (\lambda_{j,G,\boldsymbol{m}}) \in b^{\tau}_{p,q}(\mu)$, where the convergence holds unconditionally in $\mathcal{S}'(\mathbb{R}^d)$.

This definition is usually introduced as a characterization of Besov spaces. When (26) occurs, the representation is unique and we have that [44, Theorem 1.26]

$$\lambda_{j,G,\boldsymbol{m}} = 2^{jd/2}\langle f, \psi_{j,G,\boldsymbol{m}}\rangle. \tag{27}$$

To measure a given Besov regularity (fixed p, q, τ, and μ), we should select a wavelet with enough regularity that the wavelet coefficients are well-defined for $f \in B^\tau_{p,q}(\mathbb{R}^d;\mu)$. This is the meaning of (25). Under this condition, and for $f \in B^\tau_{p,q}(\mathbb{R}^d;\mu)$, the quantity

$$\|f\|_{B^\tau_{p,q}(\mathbb{R}^d;\mu)} := \left(\sum_{j\geq 0} 2^{j(\tau-d/p+d/2)q} \sum_{G\in\mathrm{G}^j} \left(\sum_{\boldsymbol{m}\in\mathbb{Z}^d} \langle 2^{-j}\boldsymbol{m}\rangle^{\mu p} |\langle f, \psi_{j,G,\boldsymbol{m}}\rangle|^p\right)^{q/p}\right)^{1/q} \tag{28}$$

is finite, with the usual modifications when p and/or $q = \infty$. The quantity (28) is a norm for $p, q \geq 1$, and a quasi-norm otherwise. In any case, the Besov space is complete for its (quasi-)norm, and is therefore a (quasi-)Banach space. We have moreover the equivalence [11, Theorem 4.2.2]

$$f \in B^\tau_{p,q}(\mathbb{R}^d;\mu) \Leftrightarrow \langle\cdot\rangle^\mu f \in B^\tau_{p,q}(\mathbb{R}^d) \tag{29}$$

with $B^\tau_{p,q}(\mathbb{R}^d) := B^\tau_{p,q}(\mathbb{R}^d;0)$ the classical (non-weighted) Besov space. The family of weighted Besov spaces includes the weighted Sobolev spaces due to the relation [11, Section 2.2.2]

$$B^\tau_{2,2}(\mathbb{R}^d;\mu) = W^\tau_2(\mathbb{R}^d;\mu). \tag{30}$$

Weighted Besov spaces are embedded, as we show in Proposition 3.

Proposition 3. *We fix $\tau_0, \tau_1, \mu_0, \mu_1 \in \mathbb{R}$ and $0 < p_0, q_0, p_1, q_1 \leq \infty$. We assume that*

$$\tau_0 > \tau_1 \text{ and } \mu_0 \geq \mu_1. \tag{31}$$

If, moreover, we have that

$$p_0 \leq p_1 \text{ and } \tau_0 - \tau_1 \geq d\left(\frac{1}{p_0} - \frac{1}{p_1}\right) \tag{32}$$

or

$$p_1 \leq p_0 \text{ and } \mu_0 - \mu_1 > d\left(\frac{1}{p_1} - \frac{1}{p_0}\right), \tag{33}$$

then we have the continuous embedding

$$B^{\tau_0}_{p_0,q_0}(\mathbb{R}^d;\mu_0) \subseteq B^{\tau_1}_{p_1,q_1}(\mathbb{R}^d;\mu_1). \tag{34}$$

Proof. Condition (32) was proved to be sufficient in [11, Section 4.2.3]. Note however that we can easily prove the embedding by using Besov sequence spaces, as we shall do for the other condition. We could not find any precise statement of embeddings between Besov spaces for $p_1 \leq p_0$ in the literature, so we provide our own proof for the sufficiency of (33).

First of all, the parameter q is dominated by parameters τ and p in the sense that, for every $\tau \geq 0$, $\epsilon > 0$, and $0 < p, q, r \leq \infty$, we have the embedding [45, Proposition 2, Section 2.3.2]

$$B^{\tau+\epsilon}_{p,q}(\mathbb{R}^d;\mu) \subseteq B^\tau_{p,r}(\mathbb{R}^d;\mu). \tag{35}$$

Note that Triebel considers unweighted spaces in [45], but the extension to the weighted case is obvious. This allows to restrict us to the case $q_0 = q_1 = q$. Fix $\boldsymbol{\lambda} = \{\lambda_{j,G,\boldsymbol{m}},\ j \geq 0, G \in$

$\mathrm{G}^j, \boldsymbol{m} \in \mathbb{Z}^d\}$. Due to the Hölder inequality, as soon as $1/a + 1/b = 1$, we have, for every $j \geq 0$ and $G \in \mathrm{G}^j$, that

$$\sum_{\boldsymbol{m}\in\mathbb{Z}^d} \langle 2^{-j}\boldsymbol{m}\rangle^{\mu_1 p_1} |\lambda_{j,G,\boldsymbol{m}}|^{p_1} \leq \left(\sum_{\boldsymbol{m}\in\mathbb{Z}^d} \langle 2^{-j}\boldsymbol{m}\rangle^{(\mu_1-\mu_0)p_1 b} \right)^{1/b} \left(\sum_{\boldsymbol{m}\in\mathbb{Z}^d} \langle 2^{-j}\boldsymbol{m}\rangle^{\mu_0 p_1 a} |\lambda_{j,G,\boldsymbol{m}}|^{p_1 a} \right)^{1/a}. \tag{36}$$

We choose $a = p_0/p_1 \geq 1$, thus $(\mu_1 - \mu_0)p_1 b = (\mu_1 - \mu_0)/(1/p_1 - 1/p_0) < -d$ by using (33), and $\sum_{\boldsymbol{m}\in\mathbb{Z}^d} \langle 2^{-j}\boldsymbol{m}\rangle^{(\mu_1-\mu_0)p_1 b} < \infty$. Since $ap_1 = p_0$, we can rewrite (36) as

$$\left(\sum_{\boldsymbol{m}\in\mathbb{Z}^d} \langle 2^{-j}\boldsymbol{m}\rangle^{\mu_1 p_1} |\lambda_{j,G,\boldsymbol{m}}|^{p_1} \right)^{1/p_1} \leq C \left(\sum_{\boldsymbol{m}\in\mathbb{Z}^d} \langle 2^{-j}\boldsymbol{m}\rangle^{\mu_0 p_0} |\lambda_{j,G,\boldsymbol{m}}|^{p_0} \right)^{1/p_0} \tag{37}$$

with $C > 0$ a finite constant. Using (24), this implies that $\|\boldsymbol{\lambda}\|_{b^{\tau_1}_{p_1,q}(\mu_1)} \leq C' \|\boldsymbol{\lambda}\|_{b^{\tau_0}_{p_0,q}(\mu_0)}$ and, then, the corresponding embedding between Besov sequence spaces. Finally, (34) is a consequence of the isomorphism between Besov sequence spaces and Besov function spaces in Definition 4 (see [44, Theorem 1.26] for more details on the isomorphism). We let the reader adapt the proof when p and/or q are infinite. □

If the only knowledge provided to us is that the generalized function f is in $\mathcal{S}'(\mathbb{R}^d)$, then this is not enough to of which regularity u should be the wavelet to characterize the Besov smoothness of f. But if we have additional information on f, for instance its inclusion in a Sobolev space, then the situation is different. Proposition gives a wavelet-domain criterion to determine if a generalized function f, known to be in $W_2^{\tau_0}(\mathbb{R}^d;\mu_0)$, is actually in a given Besov space $B^{\tau}_{p,q}(\mathbb{R}^d;\mu)$.

Proposition 4. *Let $\tau, \tau_0, \mu, \mu_0 \in \mathbb{R}$ and $0 < p, q \leq \infty$. We set*

$$u > \max(|\tau_0|, |\tau - d(1/p - 1/2)_+|). \tag{38}$$

Then, the generalized function $f \in W_2^{\tau_0}(\mathbb{R};\mu_0)$ is in $B^{\tau}_{p,q}(\mathbb{R}^d;\mu)$ if and only if

$$\sum_{j\geq 0} 2^{j(\tau - d/p + d/2)q} \sum_{G\in\mathrm{G}^j} \left(\sum_{\boldsymbol{m}\in\mathbb{Z}^d} \langle 2^{-j}\boldsymbol{m}\rangle^{\mu p} |\langle f, \psi_{j,G,\boldsymbol{m}}\rangle|^p \right)^{q/p} < \infty, \tag{39}$$

with $(\psi_{j,G,\boldsymbol{m}})$ a wavelet basis of $L_2(\mathbb{R}^d)$ of regularity u, with the usual modifications when p and/or $q = \infty$.

Proof. Let $\tau_1 < \min(\tau_0, \tau - d(1/p - 1/2)_+)$ and $\mu_1 \leq \min(\mu_0, \mu - d(1/p - 1/2)_+)$. Then, according to Proposition 3, we have the embeddings

$$B^{\tau}_{p,q}(\mathbb{R}^d;\mu) \subseteq W_2^{\tau_1}(\mathbb{R}^d;\mu_1) \text{ and } W_2^{\tau_0}(\mathbb{R}^d;\mu_0) \subseteq W_2^{\tau_1}(\mathbb{R}^d;\mu_1).$$

Condition (38) implies that we can apply Definition 4 into the space $W_2^{\tau_1}(\mathbb{R}^d;\mu_1)$. In particular, if $(\psi_{j,G,\boldsymbol{m}})$ is the wavelet basis of Definition 4 with regularity u, and for every function $f \in W_2^{\tau_1}(\mathbb{R}^d;\mu_1)$, then the wavelet coefficients $\langle f, \psi_{j,G,\boldsymbol{m}}\rangle$ are well-defined. Moreover, we have the characterization

$$f \in B^{\tau}_{p,q}(\mathbb{R}^d;\mu) \Leftrightarrow \|f\|_{B^{\tau}_{p,q}(\mathbb{R}^d;\mu)} < \infty$$

for $f \in W_2^{\tau_1}(\mathbb{R}^d;\mu_1)$ and, therefore, for $f \in W_2^{\tau_1}(\mathbb{R}^d;\mu_0)$. □

3 Moment Estimates for Lévy White Noises

The goal of this section is to obtain bounds for the pth moments of the random variable $\langle w, \varphi \rangle$, where w is a Lévy white noise and $\varphi \in \mathcal{S}(\mathbb{R}^d)$. The bounds are related to the moments of φ. For instance, for a symmetric α-stable white noise w_α, it is known [13, Lemma 2] that

$$\mathbb{E}\left[|\langle w_\alpha, \varphi \rangle|^p\right] = C_{p,\alpha} \|\varphi\|_\alpha^p, \tag{40}$$

where $C_{p,\alpha}$ is a constant, finite if and only if $\alpha = 2$ (Gaussian case), or $p < \alpha < 2$ (non-Gaussian case).

3.1 Blumenthal-Getoor Indices

In order to generalize (40) to non-stable white noises, we consider the Blumenthal-Getoor indices introduced in [4], which are classical tools to estimate the moments of Lévy processes [10, 24, 27].

Definition 5. Let f be a Lévy exponent. We consider the two intervals

$$I_0 = \left\{ p \in [0,2], \quad \limsup_{|\xi| \to 0} \frac{f(\xi)}{|\xi|^p} < \infty \right\}, \tag{41}$$

$$I_\infty = \left\{ p \in [0,2], \quad \limsup_{|\xi| \to \infty} \frac{f(\xi)}{|\xi|^p} < \infty \right\}. \tag{42}$$

The *indices of Blumenthal-Getoor* are defined by

$$\beta_0 = \sup I_0, \quad \beta_\infty = \inf I_\infty. \tag{43}$$

Proposition 5. *Consider a Lévy exponent f with intervals I_0 and I_∞ as in (41) and (42). Then, for $\tilde{\beta}_0 \in I_0$ and $\tilde{\beta}_\infty \in I_\infty$, we have the inequality*

$$\int_{\mathbb{R}^d} |f(\varphi(\boldsymbol{x}))| \mathrm{d}\boldsymbol{x} \leq C \left(\|\varphi\|_{\tilde{\beta}_0}^{\tilde{\beta}_0} + \|\varphi\|_{\tilde{\beta}_\infty}^{\tilde{\beta}_\infty} \right) \tag{44}$$

for all $\varphi \in L_{\tilde{\beta}_0}(\mathbb{R}^d) \bigcap L_{\tilde{\beta}_\infty}(\mathbb{R}^d)$ and some constant $C > 0$.

Proof. The functions $\xi \mapsto |f(\xi)|$ and $\xi \mapsto |\xi|^{\tilde{\beta}_0} + |\xi|^{\tilde{\beta}_\infty}$ are two continuous functions, the second one dominating the first one at zero and at infinity. Therefore, there exists a constant $C > 0$ that satisfies

$$|f(\xi)| \leq C \left(|\xi|^{\tilde{\beta}_0} + |\xi|^{\tilde{\beta}_\infty} \right).$$

Integrating the latter equation over $\boldsymbol{x} \in \mathbb{R}^d$ with $\xi = \varphi(\boldsymbol{x})$, we obtain (44). □

3.2 Moment Estimates for $\langle w, \varphi \rangle$

We estimate the moments of a random variable by relaying the fractional moments to the characteristic function. Proposition 6 can be found for instance in [10, 26, 28] with some variations. For the sake of completeness, we recall the proof, similar to the one of [10].

Proposition 6. *For a random variable X with characteristic function Φ_X and $0 < p < 2$, we have the relation*

$$\mathbb{E}[|X|^p] = c_p \int_{\mathbb{R}} \frac{1 - \Re(\Phi_X)(\xi)}{|\xi|^{p+1}} \mathrm{d}\xi \in [0, \infty], \tag{45}$$

for some finite constant $c_p > 0$, where $\Re(z)$ denotes the real part of $z \in \mathbb{C}$.

Proof. For $p \in (0,2)$, we have, for every $x \in \mathbb{R}$,

$$h(x) = \int_{\mathbb{R}} (1 - \cos(x\xi)) \frac{\mathrm{d}\xi}{|\xi|^{p+1}} = \left(\int_{\mathbb{R}} (1 - \cos(u)) \frac{\mathrm{d}u}{|u|^{p+1}} \right) |x|^p, \tag{46}$$

which is obtained by the change of variable $u = x\xi$. Applying this relation to $x = X$, we have, by Fubini's theorem and denoting $c_p = \left(\int_{\mathbb{R}} (1 - \cos(u)) \frac{\mathrm{d}u}{|u|^{p+1}} \right)^{-1}$, that

$$\mathbb{E}[|X|^p] = c_p \mathbb{E}\left[\int_{\mathbb{R}} (1 - \cos(\xi X)) \frac{\mathrm{d}\xi}{|\xi|^{p+1}} \right] \tag{47}$$

$$= c_p \int_{\mathbb{R}} (1 - \Re(\mathbb{E}[\mathrm{e}^{\mathrm{i}\xi X}])) \frac{\mathrm{d}\xi}{|\xi|^{p+1}} \tag{48}$$

$$= c_p \int_{\mathbb{R}} \frac{1 - \Re(\Phi_X)(\xi)}{|\xi|^{p+1}} \mathrm{d}\xi. \tag{49}$$

□

Theorem 2. *Consider a Lévy white noise w with Blumenthal-Getoor indices β_0 and β_∞. Then, for every $\varphi \in \mathcal{S}(\mathbb{R}^d)$ and $0 < p < \beta_0$, we have the inequality*

$$\mathbb{E}\left[|\langle w, \varphi \rangle|^p \right] \leq C \left(\|\varphi\|_{\tilde{\beta}_0}^p + \|\varphi\|_{\tilde{\beta}_\infty}^p \right) \tag{50}$$

for some constant $C > 0$, with $\tilde{\beta}_0 \in I_0$, $\tilde{\beta}_\infty \in I_\infty$, and $p < \tilde{\beta}_0, \tilde{\beta}_\infty$. Moreover, the result is still valid for $p = \tilde{\beta}_0 = \tilde{\beta}_\infty = 2$ if $\beta_0 = 2 \in I_0$ (finite-variance case).

We use three simple lemmas in the proof of Theorem 2.

Lemma 1. *There exists a constant $C > 0$ such that, for every $z \in \mathbb{C}$ with $\Re(z) \leq 0$, we have that*

$$|1 - \mathrm{e}^z| \leq C \left(1 - \mathrm{e}^{-|z|} \right). \tag{51}$$

Proof. First, for any $z \in \mathbb{C}$, we have that $(1 - \mathrm{e}^{-|z|}) \geq 0$. Consider the function

$$h(z) = \frac{|1 - \mathrm{e}^z|}{1 - \mathrm{e}^{-|z|}}, \tag{52}$$

which is well-defined and continuous over $\mathbb{C}\backslash\{0\}$. Since $h(z) \to 1$ when $z \to 0$, we extend h continuously to $\mathbb{C}$ with $h(0) = 1$. For z such that $\Re(z) \leq 0$ and $|z| \geq \log 2$, we have $(1 - \mathrm{e}^{-|z|}) \geq 1/2$ and $|1 - \mathrm{e}^z| \leq 2$, hence $0 \leq h(z) \leq 4$. Moreover,

$$K = \{ z \in \mathbb{C}, \quad |z| \leq \log 2, \ \Re(z) \leq 0 \}$$

is compact. Therefore, the continuous function h is bounded on K. Finally, we have shown that h is bounded on $\{z \in \mathbb{C}, \Re(z) \leq 0\}$, which gives (51). □

Lemma 2. *Let $x, y \leq 1$. Then, $(1 - xy) \leq (1 - x) + (1 - y)$.*

Proof. From the assumption, we know that $0 \leq (1 - x)(1 - y) = (1 - x) + (1 - y) - (1 - xy)$ and the result follows. □

Lemma 3. *For every $\alpha \in (0,2)$ and $p < \alpha$, there exists a constant $c_{p,\alpha}$ such that*

$$\int_{\mathbb{R}} \frac{1 - \mathrm{e}^{-|x\xi|^\alpha}}{|\xi|^{p+1}} \mathrm{d}\xi = c_{p,\alpha} |x|^p \tag{53}$$

for every $x \in \mathbb{R}$.

Proof. The result is deduced by the change of variable $u = x\xi$ in the integral. □

Proof of Theorem 2. Defining $X = \langle w, \varphi \rangle$, the characteristic function of X is

$$\Phi_X(\xi) = \exp\left(\int_{\mathbb{R}^d} f(\xi \varphi(\boldsymbol{x})) \mathrm{d}\boldsymbol{x} \right). \tag{54}$$

Moreover, from Proposition 5, we have that

$$\int_{\mathbb{R}^d} |f(\xi\varphi)| \leq C \left(\|\varphi\|_{\tilde{\beta}_0}^{\tilde{\beta}_0} |\xi|^{\tilde{\beta}_0} + \|\varphi\|_{\tilde{\beta}_\infty}^{\tilde{\beta}_\infty} |\xi|^{\tilde{\beta}_\infty} \right). \tag{55}$$

We have, therefore, that

$$\begin{aligned}
1 - \Re(\Phi_X)(\xi) &\leq |1 - \Phi_X(\xi)| \\
&\overset{(i)}{\leq} C \left(1 - \exp\left(- \left| \int f(\xi\varphi) \right| \right) \right) \\
&\overset{(ii)}{\leq} C \left(1 - \exp\left(- \int |f(\xi\varphi)| \right) \right) \\
&\overset{(iii)}{\leq} C' \left(1 - \exp(-\|\varphi\|_{\tilde{\beta}_0}^{\tilde{\beta}_0} |\xi|^{\tilde{\beta}_0}) \exp(-\|\varphi\|_{\tilde{\beta}_\infty}^{\tilde{\beta}_\infty} |\xi|^{\tilde{\beta}_\infty}) \right) \\
&\overset{(iv)}{\leq} C' \left(\left(1 - \exp(-\|\varphi\|_{\tilde{\beta}_0}^{\tilde{\beta}_0} |\xi|^{\tilde{\beta}_0}) \right) + \left(1 - \exp(-\|\varphi\|_{\tilde{\beta}_\infty}^{\tilde{\beta}_\infty} |\xi|^{\tilde{\beta}_\infty}) \right) \right),
\end{aligned} \tag{56}$$

where (i) comes from Lemma 1, (ii) and (iii) from the fact that $x \mapsto 1 - \mathrm{e}^{-x}$ is increasing, (iii) from (55), and (iv) from Lemma 2. Finally, by applying Lemma 3 with $x = \|\varphi\|_{\tilde{\beta}_0}, \alpha = \tilde{\beta}_0$ and $x = \|\varphi\|_{\tilde{\beta}_\infty}, \alpha = \tilde{\beta}_\infty$, respectively, we deduce using (45) that

$$\mathbb{E}[|X|^p] = c_p \int_{\mathbb{R}} \frac{1 - \Re(\Phi_X)(\xi)}{|\xi|^{p+1}} \mathrm{d}\xi \leq C'' \left(\|\varphi\|_{\tilde{\beta}_0}^p + \|\varphi\|_{\tilde{\beta}_\infty}^p \right), \tag{57}$$

ending the proof.

The finite-variance case, when $\beta_0 = 2 \in I_0$, cannot be deduced with the same arguments, since (45) is not valid any more. However, we know in this case that

$$\mathbb{E}[\langle w, \varphi \rangle^2] = \sigma^2 \|\varphi\|_2^2 + \mu^2 \left(\int_{\mathbb{R}^d} \varphi \right)^2 \leq \sigma^2 \|\varphi\|_2^2 + \mu^2 \|\varphi\|_1^2, \tag{58}$$

where σ^2 and μ are the variance and the mean of the infinitely divisible random variable with the same Lévy exponent as w [47, Proposition 4.15], respectively. Hence, the result is still valid. □

We take advantage of Theorem 2 in a slightly less general form and apply it to wavelets, which are functions that are rescaled versions of an initial function.Corollary 1 is a useful consequence of Theorem 2. For $\varphi \in \mathcal{S}(\mathbb{R}^d)$, $j \geq 0$, and $\boldsymbol{m} \in \mathbb{Z}^d$, we set $\varphi_{j,k} = 2^{jd/2} \varphi(2^j \cdot - \boldsymbol{m})$.

Corollary 1. *Let w be a Lévy white noise with indices β_0 and β_∞. We assume either that $\beta_\infty < \beta_0$, or that $\beta_\infty = \beta_0 \in I_\infty \bigcap I_0$. We fix $p < \beta \in I_0 \bigcap I_\infty$. Then, there exists a constant C such that, for every $\varphi \in \mathcal{S}(\mathbb{R}^d)$, $j \geq 0$, and $\boldsymbol{m} \in \mathbb{Z}^d$,*

$$\mathbb{E}\left[|\langle w, \varphi_{j,\boldsymbol{m}}\rangle|^p\right] \leq C 2^{jdp(1/2-1/\beta)} \|\varphi\|_\beta^p. \tag{59}$$

Moreover, this result is still valid if $p = \beta = 2 \in I_0$.

Proof. Remark first that the assumptions on β_0 and β_∞ imply that $I_0 \bigcap I_\infty \neq \emptyset$. We apply Theorem 2 with $\tilde{\beta}_\infty = \tilde{\beta}_0 = \beta$. In particular, we have that $\mathbb{E}\left[|\langle w, \varphi_{j,\boldsymbol{m}}\rangle|^p\right] \leq C\|\varphi_{j,\boldsymbol{m}}\|_\beta^p$. The result follows from the relation

$$\|\varphi_{j,\boldsymbol{m}}\|_\beta^p = 2^{jdp/2}\left(\int_{\mathbb{R}^d} |\varphi(2^j\boldsymbol{x} - \boldsymbol{m})|^\beta \mathrm{d}\boldsymbol{x}\right)^{p/\beta} = 2^{jdp(1/2-1/\beta)}\|\varphi\|_\beta^p, \tag{60}$$

the last equality being obtained by the change of variable $\boldsymbol{y} = 2^j\boldsymbol{x} - \boldsymbol{m}$. The result is still valid for $p = \beta = 2$ for which we can still apply Theorem 2. □

3.3 Application of Moment Estimates to the Extension of $\langle w, \varphi\rangle$ for Non-Smooth Functions

A generalized random process s is a random variable from Ω to $\mathcal{S}'(\mathbb{R}^d)$. Alternatively, it can be seen as a linear and continuous map from $\mathcal{S}(\mathbb{R}^d)$ to the space $L_0(\Omega)$ of real random variables, that associates to $\varphi \in \mathcal{S}(\mathbb{R}^d)$ the random variable $\langle s, \varphi\rangle$[1]. The space $L_0(\Omega)$ is a Fréchet space associated with the following notion of convergence. A sequence (X_n) converges to X in $L_0(\Omega)$ if and only if $\|X_n - X\|_{L_0(\Omega)} = \mathbb{E}[\min(|X_n - X|, 1)] \underset{n\to\infty}{\longrightarrow} 0$. The topology on $L_0(\Omega)$ corresponds to the convergence in probability. We also define the spaces $L_p(\Omega)$ for $0 < p < \infty$ associated for $p \geq 1$ ($p < 1$, respectively) with the norm (the quasi-norm, respectively) $\|X\|_{L_p(\Omega)} = \mathbb{E}[|X|^p]$. See [22, Section 2.2] for more details.

To measure the Besov regularity of Lévy white noises, we shall consider random variables $\langle w, \varphi\rangle$ for test functions φ not in $\mathcal{S}(\mathbb{R}^d)$. We handle this by extending the domain of test functions through which one can observe a generalized random process.

Lemma 4. *Let $0 < p, \beta < \infty$. Consider a generalized random process s. We assume that, for all $\varphi \in \mathcal{S}(\mathbb{R}^d)$,*

$$\mathbb{E}[|\langle s, \varphi\rangle|^p] \leq C\|\varphi\|_\beta^p \tag{61}$$

for some constant $C > 0$. Then, we can extend s as a linear and continuous map from $L_\beta(\mathbb{R}^d)$ to $L_0(\Omega)$. Moreover, (61) remains valid for $\varphi \in L_\beta(\mathbb{R}^d)$.

Before proving this result, we remark that it immediately implies Corollary 2.

Corollary 2. *Under the conditions of Corollary 1, we can extend $\langle w, \varphi\rangle$ for $\varphi \in L_\beta(\mathbb{R}^d)$. Moreover, (59) remains valid for any $\varphi \in L_\beta(\mathbb{R}^d)$.*

[1] This is not as obvious as it might seem in infinite-dimension, and is again due to the nuclear structure of $\mathcal{S}(\mathbb{R}^d)$. For the links between E'-valued random variables and linear functionals from E to $L_0(\Omega)$ (with E' the dual of E), see [22, Section 2.3], in particular Theorems 2.3.1 and 2.3.2.

Proof of Lemma 4. Fix $\varphi \in L_\beta(\mathbb{R}^d)$. Let (φ_n) be a sequence of elements in $\mathcal{S}(\mathbb{R}^d)$ such that $\varphi_n \underset{n\to\infty}{\longrightarrow} \varphi$ in $L_\beta(\mathbb{R}^d)$. Such a sequence exists by density of $\mathcal{S}(\mathbb{R}^d)$ into $L_\beta(\mathbb{R}^d)$, well-known for $\beta \geq 1$, and easily extended for $0 < \beta < 1$. Then, for every $n, m \in \mathbb{N}$, it holds that

$$\begin{aligned}\mathbb{E}[|\langle s, \varphi_n\rangle - \langle s, \varphi_m\rangle|^p] &= \mathbb{E}[|\langle s, \varphi_n - \varphi_m\rangle|^p] \\ &\leq C\|\varphi_n - \varphi_m\|_\beta^p,\end{aligned}$$

so that $(\langle s, \varphi_n\rangle)$ is a Cauchy sequence in the complete space $L_p(\Omega)$. It therefore converges to a random variable X in $L_p(\Omega)$. Moreover, this limit does not depend on the sequence (φ_n) chosen to approximate φ. Indeed, if $(\tilde{\varphi}_n)$ is another sequence converging to φ in $L_\beta(\mathbb{R}^d)$, and defining $\tilde{X}$ as the limit of $\langle s, \tilde{\varphi}_n\rangle$ in $L_p(\Omega)$, we have that

$$\mathbb{E}[|X - \tilde{X}|^p] = \mathbb{E}[|X - \langle s, \varphi_n\rangle + \langle s, \varphi_n - \tilde{\varphi}_n\rangle + \langle s, \tilde{\varphi}_n\rangle - \tilde{X}|^p] \tag{62}$$

We have moreover, for $x, y, z \in \mathbb{R}$, that

$$|x + y + z|^p \leq \begin{cases} 3^{p-1}\left(|x|^p + |y|^p + |z|^p\right) & \text{if } p \geq 1, \\ |x|^p + |y|^p + |z|^p & \text{otherwise.}\end{cases}$$

Hence, if we set $C_p = 3^{p-1}$ if $p \geq 1$ and $C_p = 1$ if $p < 1$, we deduce from (62) that

$$\mathbb{E}[|X - \tilde{X}|^p] \leq C_p\left(\mathbb{E}[|X - \langle s, \varphi_n\rangle|^p] + \mathbb{E}[\langle s, \varphi_n - \tilde{\varphi}_n\rangle|^p] + \mathbb{E}[\langle s, \tilde{\varphi}_n\rangle - \tilde{X}|^p]\right). \tag{63}$$

The three terms in (63) vanish when $n \to \infty$. For the second one, we use that $\mathbb{E}[\langle s, \varphi_n - \tilde{\varphi}_n\rangle|^p] \leq C\|\varphi_n - \tilde{\varphi}_n\|_\beta^p$. This proves that $X = \tilde{X}$ almost surely. We therefore define $X = \langle s, \varphi\rangle$.

Finally, (61) is still valid for $\varphi \in L_\beta(\mathbb{R}^d)$ by continuity of s from $L_\beta(\mathbb{R}^d)$ to $L_p(\Omega)$. □

4 Measurability of Weighted Besov Spaces

A generalized random process is a measurable function from Ω to $\mathcal{S}'(\mathbb{R}^d)$, endowed with the cylindrical σ-field $\mathcal{B}_c(\mathcal{S}'(\mathbb{R}^d))$. In the next sections, we investigate in which Besov space (local or weighted) is a given Lévy white noise. We show here that this question is meaningful in the sense that any Besov space $B^\tau_{p,q}(\mathbb{R}^d; \mu)$ is measurable in $\mathcal{S}'(\mathbb{R}^d)$.

Proposition 7. *For every $0 < p, q \leq \infty$ and $\tau, \mu \in \mathbb{R}$, we have that*

$$B^\tau_{p,q}(\mathbb{R}^d; \mu) \in \mathcal{B}_c(\mathcal{S}'(\mathbb{R}^d)). \tag{64}$$

The proof of this result is very similar to the one of [13, Theorem 4], except we work now over $\mathbb{R}^d$ and deal with weights. In particular, we will rely on [13, Lemma 1].

Proof. We obtain the desired result in three steps. We treat the case $p, q < \infty$ and let the reader adapt the proof for p and/or $q = \infty$.

- First, we show that $W^\tau_2(\mathbb{R}^d; \mu) \in \mathcal{B}_c(\mathcal{S}'(\mathbb{R}^d))$ for every $\tau, \mu \in \mathbb{R}$. Let $(h_n)_{n\in\mathbb{N}}$ be an orthonormal basis of $L_2(\mathbb{R}^d)$, with $h_n \in \mathcal{S}(\mathbb{R}^d)$ for all $n \geq 0$. (We can for instance consider the Hermite functions, based on Hermite polynomials, see [41, Section 2] or [22, Section 1.3] for the definitions.) The interest of having basis functions in $\mathcal{S}(\mathbb{R}^d)$ is that we have the characterization

$$L_2(\mathbb{R}^d) = \left\{ f \in \mathcal{S}'(\mathbb{R}^d),\ \sum_{n\in\mathbb{N}} |\langle f, h_n\rangle|^2 < \infty \right\}. \tag{65}$$

More generally, with the notations of Section 2.2.1, $f \in W_2^\tau(\mathbb{R}^d;\mu)$ if and only if $\mathrm{L}_\tau\{\langle\cdot\rangle^\mu f\} \in L_2(\mathbb{R}^d)$, from which we deduce that

$$W_2^\tau(\mathbb{R}^d;\mu) = \left\{ f \in \mathcal{S}'(\mathbb{R}^d),\ \sum_{n\in\mathbb{N}} |\langle f, \langle\cdot\rangle^\mu \mathrm{L}_\tau\{h_n\}\rangle|^2 < \infty \right\}. \tag{66}$$

We can therefore apply [13, Lemma 1] with $\alpha = 2$, $S = \mathbb{N}$, and $\varphi_n = \langle\cdot\rangle^\mu \mathrm{L}_\tau\{h_n\}$, to deduce that $W_2^\tau(\mathbb{R}^d;\mu) \in \mathcal{B}_c(\mathcal{S}'(\mathbb{R}^d))$.

- For any $\tau, \mu \in \mathbb{R}$, the cylindrical σ-field of $W_2^\tau(\mathbb{R}^d;\mu)$ is the σ-field $\mathcal{B}_c(W_2^\tau(\mathbb{R}^d;\mu))$generated by the sets

$$\Big\{ u \in W_2^\tau(\mathbb{R}^d;\mu),\ (\langle u,\varphi_1\rangle, \cdots, \langle u,\varphi_n\rangle) \in B \Big\}, \tag{67}$$

where $N \geq 1$, $\varphi_1, \ldots, \varphi_N \in W_2^{-\tau}(\mathbb{R}^d;-\mu)$, and B is a Borelian subset of $\mathbb{R}^N$. Then, $W_2^\tau(\mathbb{R}^d;-\mu) \in \mathcal{B}_c(\mathcal{S}'(\mathbb{R}^d))$ implies that

$$\mathcal{B}_c(W_2^\tau(\mathbb{R}^d;\mu)) \subset \mathcal{B}_c(\mathcal{S}'(\mathbb{R}^d)). \tag{68}$$

- Finally, we show that $B_{p,q}^\tau(\mathbb{R}^d;\mu) \in \mathcal{B}_c(W_2^{\tau_1}(\mathbb{R}^d;\mu_1))$ for some $\tau_1, \mu_1 \in \mathbb{R}$. Coupled with (68), we deduce (64).

  Fix $\tau_1 \leq \tau + d\,(1/2 - 1/p)$ and $\mu_1 < \mu + d\,(1/p - 1/2)$. According to Proposition 3, we have the embedding $B_{p,q}^\tau(\mathbb{R}^d;\mu) \subseteq W_2^{\tau_1}(\mathbb{R}^d;\mu_1)$. Now, we can rewrite Proposition 4 (with $\tau_0 = \tau_1$ and $\mu_0 = \mu_1$) as

$$B_{p,q}^\tau(\mathbb{R}^d;\mu) = \left\{ f \in W_2^{\tau_1}(\mathbb{R}^d;\mu_1),\ \sum_{j,G} \left(\sum_{\boldsymbol{m}} |\langle f, 2^{j(\tau-d/p+d/2)} \langle 2^{-j}\boldsymbol{m}\rangle^\mu \psi_{j,G,\boldsymbol{m}}\rangle|^p \right)^{q/p} < \infty \right\}. \tag{69}$$

  Again, we apply [13, Lemma 1] with $S = \{(j,G), j \in \mathbb{Z}, G \in \mathrm{G}_j\}$, $n = (j,G)$, $T_n = T_{(j,G)} = \mathbb{Z}^d$, $\varphi_{n,m} = \varphi_{j,G,\boldsymbol{m}} = 2^{j(\tau-d/p+d/2)} \langle 2^{-j}\boldsymbol{m}\rangle^\mu \psi_{j,G,\boldsymbol{m}}$, $\alpha = p$, and $\beta = q/p$, to deduce that $B_{p,q}^\tau(\mathbb{R}^d;\mu) \in \mathcal{B}_c(W_2^{\tau_1}(\mathbb{R}^d;\mu_1))$. Remark that, strictly speaking, Lemma 1 in [13] is stated for T_n finite, but the proof is easily adapted to T_n countable.

□

Proposition 7 suggests that the framework of generalized random processes is particularly well-suited to addressing regularity issues. By comparison, we recall that the space $\mathcal{C}(\mathbb{R}^d)$ of continuous functions is not measurable with respect to the topological σ-field on the space of all functions, while we have that

$$\mathcal{C}(\mathbb{R}^d) \in \mathcal{B}_c(\mathcal{D}'(\mathbb{R}^d)), \tag{70}$$

which is the cylindrical σ-field of the space of generalized functions (not necessarily tempered) [14, Proposition III.3.3]. See [7] for a discussion on the measurability of function spaces and the advantages of generalized random processes.

5 Lévy White Noises on Weighted Sobolev Spaces

In order to characterize the Besov smoothness of Lévy white noises, we first obtain information on their Sobolev smoothness.

Proposition 8. *A Lévy white noise w with indices $\beta_0 > 0$ and β_∞ is in the weighted Sobolev space $W_2^{-\tau}(\mathbb{R}^d; -\mu)$ if*

$$\mu > \frac{d}{\beta_0} \text{ and } \tau > \frac{d}{2}. \tag{71}$$

Proof. As we have seen in Proposition 1, we have the countable projective limit

$$\mathcal{S}(\mathbb{R}^d) = \bigcap_{\tau,\mu\in\mathbb{R}} W_2^\tau(\mathbb{R}^d;\mu) = \bigcap_{n\in\mathbb{N}} W_2^n(\mathbb{R}^d;n).$$

We are therefore in the context of [19, Theorem A.2]. It implies in particular that, if

- the characteristic functional $\widehat{\mathscr{P}}_w$ of w is continuous over $L_2(\mathbb{R}^d;\mu_0)$, and
- the identity operator I is Hilbert-Schmidt from $W_2^\tau(\mathbb{R}^d;\mu)$ to $L_2(\mathbb{R}^d;\mu_0)$,

then $w \in W_2^{-\tau}(\mathbb{R}^d, -\mu) = \left(W_2^\tau(\mathbb{R}^d;\mu)\right)'$ almost surely.

Continuity of $\widehat{\mathscr{P}}_w$ Let $\mu_0 > d\left(\frac{1}{\beta_0} - \frac{1}{2}\right) \geq 0$. Fix $\epsilon > 0$ small enough such that

$$\beta_0 - \epsilon > 0 \text{ and } \mu_0 > d\left(\frac{1}{\beta_0 - \epsilon} - \frac{1}{2}\right). \tag{72}$$

Applying Proposition 5 with $\tilde{\beta}_\infty = 2$ and $\tilde{\beta}_0 = \beta_0 - \epsilon$, we deduce that

$$\int_{\mathbb{R}^d} |f(\varphi)| \leq C\left(\|\varphi\|_{\tilde{\beta}_0}^{\tilde{\beta}_0} + \|\varphi\|_2^2\right). \tag{73}$$

Since $\mu_0 > 0$, we have that $\|\varphi\|^2_{L_2(\mathbb{R}^d)} \leq \|\varphi\|^2_{L_2(\mathbb{R}^d;\mu_0)}$. Moreover, using Hölder inequality, we have

$$\|\varphi\|_{\tilde{\beta}_0}^{\tilde{\beta}_0} = \int |\varphi|^{\tilde{\beta}_0} \leq \int (|\varphi|^{\tilde{\beta}_0} \langle\cdot\rangle^{\mu_0\tilde{\beta}_0})^p \int \langle\cdot\rangle^{-\mu_0\tilde{\beta}_0 q} \tag{74}$$

for any $1 \leq p, q \leq \infty$ such that $1/p + 1/q = 1$. Setting $p = 2/\tilde{\beta}_0 \geq 1$, we have $q = \frac{2}{2-\tilde{\beta}_0}$. Therefore,

$$\|\varphi\|_{\tilde{\beta}_0}^{\tilde{\beta}_0} \leq \int (|\varphi|\langle\cdot\rangle^{\mu_0})^2 \int \langle\cdot\rangle^{-\frac{2\tilde{\beta}_0\mu_0}{2-\tilde{\beta}_0}}, \tag{75}$$

the last integral being finite due to (72), which implies that $\frac{2\tilde{\beta}_0\mu_0}{2-\tilde{\beta}_0} > d$. Finally, injecting (75) into (73), we obtain the inequalities

$$|\log \widehat{\mathscr{P}}_w(\varphi)| \leq \int_{\mathbb{R}^d} |f(\varphi)| \leq C'\|\varphi\|^2_{L_2(\mathbb{R}^d;-\mu_0)}. \tag{76}$$

This implies that $\widehat{\mathscr{P}}_w$ is well-defined over $L_2(\mathbb{R}^d;-\mu_0)$ and continuous at $\varphi = 0$. Since $\widehat{\mathscr{P}}_w$ is positive-definite, it is therefore continuous over $L_2(\mathbb{R}^d;-\mu_0)$ [20].

Hilbert-Schmidt condition The operator $\mathrm{L}_{-\tau,-\mu}$ is defined in Proposition 1. Its kernel $k_{-\tau,-\mu}$, such that

$$\mathrm{L}_{-\tau,-\mu}\{\varphi\}(\boldsymbol{x}) = \int_{\mathbb{R}^d} k_{-\tau,-\mu}(\boldsymbol{x},\boldsymbol{y})\varphi(\boldsymbol{y})\mathrm{d}\boldsymbol{y}, \tag{77}$$

is given by

$$k_{-\tau,-\mu}(\boldsymbol{x},\boldsymbol{y}) = \langle \boldsymbol{x}\rangle^{-\mu} k_{-\tau}(\boldsymbol{y}-\boldsymbol{x}) \tag{78}$$

with $k_{-\tau}$ being the Fourier multiplier of $\mathrm{L}_{-\tau}$, which satisfies $\widehat{k}_{-\tau}(\boldsymbol{\omega}) = \langle \boldsymbol{\omega}\rangle^{-\tau}$.

According to Proposition 1, the operator $\mathrm{L}_{-\tau,-\mu+\mu_0}$ is an isometry from $L_2(\mathbb{R}^d)$ to $W_2^\tau(\mathbb{R}^d;\mu-\mu_0)$. Therefore, we have the equivalences

$$\begin{aligned}\mathrm{I} : W_2^\tau(\mathbb{R}^d;\mu) \to L_2(\mathbb{R}^d;\mu_0) \text{ is HS} &\Leftrightarrow \mathrm{I} : W_2^\tau(\mathbb{R}^d;\mu-\mu_0) \to L_2(\mathbb{R}^d;\mu_0) \text{ is HS} \\ &\Leftrightarrow \mathrm{L}_{-\tau,-\mu+\mu_0} : L_2(\mathbb{R}^d) \to L_2(\mathbb{R}^d) \text{ is HS.}\end{aligned} \tag{79}$$

There HS means Hilbert-Schmidt. The last condition in (79) is equivalent with the condition [31, Theorem VI.23]

$$k_{-\tau,-\mu+\mu_0}(\cdot,\cdot) \in L_2(\mathbb{R}^d \times \mathbb{R}^d). \tag{80}$$

Moreover, by the change of variable $\boldsymbol{z} = (\boldsymbol{y}-\boldsymbol{x})$ and using Parseval relation,

$$\begin{aligned}\int_{\mathbb{R}^d\times\mathbb{R}^d} k_{-\tau,-\mu+\mu_0}(\boldsymbol{x},\boldsymbol{y})^2 \mathrm{d}\boldsymbol{x}\mathrm{d}\boldsymbol{y} &= \int_{\mathbb{R}^d} \langle \boldsymbol{x}\rangle^{-2(\mu-\mu_0)}\mathrm{d}\boldsymbol{x} \int_{\mathbb{R}^d} k_{-\tau}(\boldsymbol{z})\mathrm{d}\boldsymbol{z} &(81)\\ &= \|\langle\cdot\rangle^{-(\mu-\mu_0)}\|^2_{L_2(\mathbb{R}^d)} \|\langle\cdot\rangle^{-\tau}\|^2_{L_2(\mathbb{R}^d)}. &(82)\end{aligned}$$

Therefore, $k_{\tau,\mu-\mu_0}$ is square-integrable if and only if $(\mu-\mu_0) > d/2$ and $\tau > d/2$. By choosing μ_0 close to $d(1/\beta_0 - 1/2)$, we finally obtain that $w \in W_2^{-\tau}(\mathbb{R}^d;-\mu)$ if

$$\tau > \frac{d}{2} \text{ and } \mu > d\left(\frac{1}{\beta_0}-\frac{1}{2}\right)+\frac{d}{2} = \frac{d}{\beta_0}, \tag{83}$$

and (71) is proved. □

6 Lévy White Noises on Weighted Besov Spaces

We investigate here the Besov smoothness of Lévy white noises over the complete space $\mathbb{R}^d$. The paths of a nontrivial white noise w are never included in $B^\tau_{p,q}(\mathbb{R}^d)$, since their is no decay at infinity. For this reason, and as for Sobolev spaces, we consider the weighted Besov spaces, introduced in Section 2.2. The main course of this section is to prove Theorem 3. .

Theorem 3. *Consider a Lévy white noise w with Blumenthal-Getoor indices $\beta_0 > 0$ and β_∞. Let $0 < p,q \le \infty$, $\tau,\mu \in \mathbb{R}$. If*

$$\mu > \frac{d}{\min(p,\beta_0)} \text{ and } \tau > d\left(1-\frac{1}{\max(p,\beta_\infty)}\right), \tag{84}$$

then $w \in B^{-\tau}_{p,q}(\mathbb{R}^d;-\mu)$ a.s.

Proof. We start with some preliminary remarks.

- First of all, it is sufficient to prove (84) for $p = q$, the other cases being deduced by the embedding relations $B^{-\tau+\epsilon}_{p,p}(\mathbb{R}^d; -\mu) \subseteq B^{-\tau}_{p,q}(\mathbb{R}^d; -\mu)$ already seen in (35). Therefore, a different parameter q can always be absorbed at the cost of an arbitrarily small smoothness, which is still possible in our case since the condition on τ in (84) is a strict inequality. For the same reason, it is admissible to consider that $p < \infty$.

- Second, we know from Proposition 8 that, for a fixed $\epsilon > 0$ and with probability 1,

$$w \in W_2^{-d/2-\epsilon}\left(\mathbb{R}^d; -\frac{d}{\beta_0} - \epsilon\right). \tag{85}$$

  From now, we fix $p = q, \tau, \mu$. We can apply Proposition 4 with $\tau_0 = -d/2 - \epsilon$ and $\mu_0 = d(\frac{1}{\beta_0} - \frac{1}{2}) + \epsilon$. We set u according to (38) and consider $(\psi_{j,G,\boldsymbol{m}})$ a wavelet basis with regularity u thereafter.

First case: $\beta_\infty < \beta_0$ or $\beta_\infty = \beta_0 \in I_0 \bigcap I_\infty$ We fix $p < \beta \in I_\infty \bigcap I_0$. We are by assumption in the conditions of Corollary 1. In particular, Corollary 2 applies: The random variable $\langle w, \varphi \rangle$ is well-defined for any $\varphi \in L_\beta(\mathbb{R}^d)$. In particular, since $\beta \in (0, 2]$, the Daubechies wavelets $\psi_{j,G,\boldsymbol{m}}$, which are compactly supported and in $L_2(\mathbb{R}^d)$, are in $L_\beta(\mathbb{R}^d)$, so that the random variables $\langle w, \psi_{j,G,\boldsymbol{m}} \rangle$ are well-defined and (59) is applicable to them. We shall show that $w \in B^{-\tau}_{p,p}(\mathbb{R}^d; -\mu)$ a.e. if

$$\mu > d/p \text{ and } \tau > d(1 - 1/\beta). \tag{86}$$

To show that $w \in B^{-\tau}_{p,p}(\mathbb{R}^d; -\mu)$ with probability 1, it is sufficient to show that

$$\mathbb{E}\left[\|w\|^p_{B^{-\tau}_{p,p}(\mathbb{R}^d;-\mu)}\right] = \sum_{j\geq 0}\left(2^{j(-\tau p - d + dp/2)} \sum_{G\in \mathrm{G}^j, \boldsymbol{m}\in\mathbb{Z}^d} \frac{\mathbb{E}[|\langle w, \psi_{j,G,\boldsymbol{m}}\rangle|^p]}{\langle 2^{-j}\boldsymbol{m}\rangle^{\mu p}}\right) < \infty. \tag{87}$$

The white noise w being stationary, $\mathbb{E}[|\langle w, \psi_{j,G,\boldsymbol{m}}\rangle|^p]$ does not depend on the shift parameter $\boldsymbol{m}$. Moreover, using (59) with $a = 2^{-j}$, we have that

$$\mathbb{E}[|\langle w, \psi_{j,G,\boldsymbol{m}}\rangle|^p] \leq C 2^{jpd(1/2-1/\beta)} \|\psi_G\|^p_\beta. \tag{88}$$

Hence, we deduce that

$$\mathbb{E}\left[\|w\|^p_{B^{-\tau}_{p,p}(\mathbb{R}^d;-\mu)}\right] \leq C' \sum_{j\geq 0} 2^{j(-\tau p - d + dp - dp/\beta)} \sum_{\boldsymbol{m}\in\mathbb{Z}^d} \langle 2^{-j}\boldsymbol{m}\rangle^{-\mu p}, \tag{89}$$

where $C' = C \sum_{G\in \mathrm{G}^0} \|\psi_G\|^p_\beta$ is a finite constant. The sum $\sum_{\boldsymbol{m}\in\mathbb{Z}^d} \langle 2^{-j}\boldsymbol{m}\rangle^{-\mu p}$ is finite if and only if $\mu > d/p$, in which case there exists a constant $C_0 > 0$ such that

$$\sum_{\boldsymbol{m}\in\mathbb{Z}^d} \langle 2^{-j}\boldsymbol{m}\rangle^{-\mu p} \underset{j\to\infty}{\sim} C_0 2^{jd}. \tag{90}$$

Indeed, we have the convergence of the Riemann sums

$$\frac{1}{n^d} \sum_{\boldsymbol{m}\in\mathbb{Z}^d} \left\langle \frac{\boldsymbol{m}}{n} \right\rangle^{-\mu p} \underset{n\to\infty}{\longrightarrow} \int_{\mathbb{R}^d} \langle \boldsymbol{x}\rangle^{-\mu p} \mathrm{d}\boldsymbol{x} < \infty \tag{91}$$

and (90) is showed setting $2^j = n$, for $C_0 = \int_{\mathbb{R}^d} \langle \boldsymbol{x}\rangle^{-\mu p} \mathrm{d}\boldsymbol{x}$. Finally, for $\mu > d/p$, the quantity $\mathbb{E}\left[\|w\|^p_{B^{-\tau}_{p,p}(\mathbb{R}^d;-\mu)}\right]$ is finite if

$$\sum_{j\geq 0} 2^{j(-\tau p + dp - dp/\beta)} < \infty, \tag{92}$$

which happens when

$$\tau - d + d/\beta > 0. \tag{93}$$

We have shown that $w \in B_{p,p}^{-\tau}(\mathbb{R}^d; -\mu)$ under the conditions of (86), as expected. We now split the domain of p:

- if $p \leq \beta_\infty$, by choosing β close enough to β_∞ (or equal if $\beta_0 = \beta_\infty$), we obtain that $w \in B_{p,p}^{-\tau}(\mathbb{R}^d; -\mu)$ if $\mu > d/p$ and $\tau > d - d/\beta_\infty$;
- if $\beta_\infty < p < \beta_0$, by choosing β close enough to p, we obtain that $w \in B_{p,p}^{-\tau}(\mathbb{R}^d; -\mu)$ if $\mu > d/p$ and $\tau > d - d/p$.

We summarize the situation by $w \in B_{p,p}^{-\tau}(\mathbb{R}^d; -\mu)$ if $\mu > d/p$ and $\tau > d - d/\max(p, \beta_\infty)$, which corresponds to (84) for $p < \beta_0$. Finally, the case $p \geq \beta_0$ is deduced from the result for $p < \beta_0$ (by considering values of p arbitrarily close to β_0) and the embedding (32).

Second case: general (β_0, β_∞) A white noise w can be decomposed as

$$w = w_1 + w_2, \tag{94}$$

where w_1 and w_2 are independent, w_1 is a compound-Poisson white noise, and w_2 is finite-variance. To see that, we invoke the Lévy-Itô decomposition, see for instance [35, Chapter 4]. It means in particular that $\beta_0(w_1) = \beta_0 > 0$ and $\beta_\infty(w_1) = 0$. Therefore, w_1 is covered by the first case. Moreover, $\beta_\infty(w_2) = \beta_\infty$ and $\beta_0(w_2) = 2 \in I_0(w_2)$. Again, w_2 is covered by the first case. Indeed, it is obvious if $\beta_\infty < 2$. But if $\beta_\infty = 2$, we have that $\beta_\infty(w_2) = \beta_0(w_2) = 2 \in I_0(w_2) \bigcap I_\infty(w_2)$. Hence, w is the sum of two processes w_1 and w_2 that are in $B_{p,q}^{-\tau}(\mathbb{R}^d; -\mu)$ under the conditions (84). Besov spaces being linear spaces, these conditions are also sufficient for w.

□

7 Lévy White Noises on Local Besov Spaces

The space of infinitely smooth and compactly supported functions is denoted by $\mathcal{D}(\mathbb{R}^d)$. Its topological dual is $\mathcal{D}'(\mathbb{R}^d)$, the space of generalized functions, not necessarily tempered. In the same way that we defined generalized random processes over $\mathcal{S}'(\mathbb{R}^d)$, we can also define generalized random processes over $\mathcal{D}'(\mathbb{R}^d)$. This is actually the original approach of Gelfand and Vilenkin in [16]. As we briefly saw in Section 2.1, the class of Lévy white noises over $\mathcal{D}'(\mathbb{R}^d)$ is strictly larger than the one over $\mathcal{S}'(\mathbb{R}^d)$. A Lévy white noise over $\mathcal{D}'(\mathbb{R}^d)$ is also in $\mathcal{S}'(\mathbb{R}^d)$ if and only if its Lévy exponent satisfies the Schwartz condition [9] or, equivalently, if and only if its Blumenthal-Getoor index β_0 is not 0. Until now, we have only considered Lévy white noises for which $\beta_0 \neq 0$. Since we shall now focus on the local Besov smoothness of Lévy white noises, the two equivalent conditions are now superfluous.

Definition 6. Let $\tau \in \mathbb{R}$ and $0 < p, q \leq \infty$. The *local Besov space* $B_{p,q}^{\tau,\mathrm{loc}}(\mathbb{R}^d)$ is the collection of functions $f \in \mathcal{D}'(\mathbb{R}^d)$ such that $f \times \varphi \in B_{p,q}^{\tau}(\mathbb{R}^d)$ for every $\varphi \in \mathcal{D}(\mathbb{R}^d)$.

The weighted and local Besov regularities are linked according to Proposition 9.

Proposition 9. *Let $\tau, \mu \in \mathbb{R}$, $0 < p, q \leq \infty$. We have the continuous embedding*

$$B_{p,q}^{\tau}(\mathbb{R}^d; \mu) \subseteq B_{p,q}^{\tau,\mathrm{loc}}(\mathbb{R}^d). \tag{95}$$

The local regularity of Lévy white noise is directly obtained from the previous results, essentially up to the case of compound-Poisson noise with $\beta_0 = 0$. Before stating the main result of this section, we therefore have to analyze the compound-Poisson case.

Definition 7. A *compound-Poisson white noise* is a Lévy white noise with a Lévy exponent of the form

$$f(\xi) = \exp\left(\lambda(\widehat{\mathcal{P}}_{\text{jump}}(\xi) - 1)\right), \tag{96}$$

where $\lambda > 0$ is called the *Poisson parameter* and $\mathcal{P}_{\text{jump}}$ is a probability law on $\mathbb{R}\backslash\{0\}$ called the *law of jumps*.

Compound-Poisson random variables are infinitely divisible [35], so that (96) defines a valid Lévy exponent.

Lemma 5. *Let $\tau \in \mathbb{R}$ and $0 < p, q \leq \infty$. Consider a compound-Poisson noise w. If*

$$\tau > d\left(1 - \frac{1}{p}\right), \tag{97}$$

then $w \in B_{p,q}^{-\tau,\text{loc}}(\mathbb{R}^d)$.

Proof. Let λ and $\mathcal{P}_{\text{jump}}$ be the Poisson parameter and the law of jumps of w, respectively. The compound-Poisson noise w can be written as [46, Theorem 1]

$$w = \sum_{n\in\mathbb{N}} a_k \delta(\cdot - \boldsymbol{x}_k), \tag{98}$$

where (a_k) are i.i.d. with law $\mathcal{P}_{\text{jump}}$ and $(\boldsymbol{x}_k)$ are such that the number of $\boldsymbol{x}_k$ in any Borelian $B \in \mathbb{R}^d$ is a Poisson random variable with parameter $\lambda\mu(B)$, μ denoting the Lebesgue measure on $\mathbb{R}^d$. For $\varphi \in \mathcal{D}(\mathbb{R}^d)$, the function φ being compactly supported, the generalized random process $w \times \varphi = \sum_{n\in\mathbb{N}} a_n \varphi(\boldsymbol{x}_n)\delta(\cdot - \boldsymbol{x}_n)$ is almost surely a finite sum of shifted Dirac functions. Hence, it has the Besov regularity of a single Dirac function, which is precisely (97); see [39, p. 164]. □

Corollary 3. *Let $0 < p, q \leq \infty$, $\tau \in \mathbb{R}$. Consider a Lévy white noise w with indices β_0, β_∞. If*

$$\tau > d\left(\frac{1}{\max(p, \beta_\infty)} - 1\right), \tag{99}$$

then $w \in B_{p,q}^{-\tau,\text{loc}}(\mathbb{R}^d)$ a.s.

Proof. The case $\beta_0 > 0$ is a direct consequence of Theorem 3 and Proposition 9. Let assume now that $\beta_0 = 0$. Again, we can split w as $w_1 + w_2$, where w_1 is compound-Poisson and w_2 is finite-variance. For w_2, we can still apply Theorem 3. We can therefore restrict our attention to the case of compound-Poisson noises with $\beta_0 = 0$. But we have seen that the compound-Poisson case—regardless of β_0— was covered in Lemma 5. Since $\beta_\infty = 0$ for compound-Poisson noises, Lemma 5 is consistent with (99), finishing the proof. □

8 Discussion and Examples

8.1 Discussion and Comparison with Known Results

Sobolev regularity of Lévy white noises It is noteworthy to observe that Sections 5 and 6, while based on very different techniques, give exactly the same results when applied to Sobolev spaces. Indeed, applying Theorem 3 with $p = q = 2$, we recover exactly (71) due to the relations $\min(2, \beta_0) = \beta_0$ and $\max(2, \beta_\infty) = 2$. Theorem 3 is therefore the generalization of Proposition 8, from Sobolev to Besov spaces.

Interestingly, the Sobolev smoothness parameter τ of a Lévy white noise does not depend on the noise: The universal sufficient condition is $\tau > d/2$. Moreover, we conjecture that this condition is also necessary, in the sense that $w \notin W_2^\tau(\mathbb{R}^d; \mu)$ with probability 1 for $\tau \geq d/2$ for any μ and any white noise w. The situation is different when considering Besov smoothness for $p \neq 2$.

Hölder regularity of Lévy white noises We obtain the Hölder regularity of a white noise by setting $p = q = \infty$ in Theorem 3. Because $\min(\infty, \beta_0) = \beta_0$ and $\max(\infty, \beta_\infty) = \infty$, we deduce Corollary 4.

Corollary 4. *The Lévy white noise w with Blumenthal-Getoor indices $\beta_0 > 0$ and β_∞ is in the weighted Hölder space $H^{-\tau}(\mathbb{R}^d; -\mu)$ if*

$$\mu > d/\beta_0, \quad \tau > d. \tag{100}$$

As for the Sobolev regularity, the Hölder regularity of a Lévy noise that we obtained is independent of the noise. However, the Gaussian white noise has a local Hölder regularity of $(-\tau)$ for every $\tau > \frac{d}{2}$ [48]. It means that our bounds for the regularity are suboptimal for the Gaussian white noise. By contrast with the Gaussian case, we conjecture that the condition $\tau > d$ is optimal for non-Gaussian Lévy white noises.

The regularity of Lévy white noises for general p Fixing the parameters $p = q > 0$, we define

$$\tau_p(w) = \min\{\tau \in \mathbb{R}, w \in B_{p,p}^{-\tau,\mathrm{loc}}(\mathbb{R}^d) \text{ a.s.}\}. \tag{101}$$

The quantity $\tau_p(w)$ measures the regularity of the Lévy white noise w for the L_p-(quasi-)norm. In Corollary 3, we have seen that $\tau_p(w) \leq d\left(\frac{1}{\max(p,\beta_\infty)} - 1\right)$, a quantity that does not depend on β_0. We conjecture that

$$\tau_p(w) = d\left(\frac{1}{\max(p, \beta_\infty)} - 1\right) \tag{102}$$

for non-Gaussian white noises. If this is true, then the quantity $\max(p, \beta_\infty)$ is a measure of the regularity of a Lévy white noise for the L_p-(quasi-)norm.

We summarize the local results of Corollary 3 with the diagram of Figure 1. We use the classical $(1/p, \tau)$-representation, which is most convenient for visualization. We indeed see in (102) that the parameters $1/p$ and τ are linked with a linear relation for $p \leq \beta_\infty$.

Weights and Lévy white noises As for the regularity, we can define for $p = q > 0$ the optimal weight

$$\mu_p(w) = \min\{\mu \in \mathbb{R},\ \exists \tau \in \mathbb{R},\ w \in B_{p,p}^{-\tau}(\mathbb{R}^d; \mu) \text{ a.s.}\}. \tag{103}$$



(a) $0 < \beta_\infty < 1$ (b) $1 < \beta_\infty < 2$

Figure 1: Besov localization of general Lévy white noises. A white noise is almost surely in a given local Besov space $B^{\tau,\mathrm{loc}}_{p,q}(\mathbb{R}^d)$ if $(1/p, \tau)$ is located in the shaded region.

Table 1: Blumental-Getoor indices of Lévy white noises

White noise	parameter	$f(\xi)$	$p_{\mathrm{id}}(x)$	β_0	β_∞	*cf.*
Gaussian	$\sigma^2 > 0$	$-\sigma^2\xi^2/2$	$\frac{\mathrm{e}^{-x^2/2\sigma^2}}{\sqrt{2\pi\sigma^2}}$	2	2	[48]
Pure drift	$\mu \in \mathbb{R}$	$\mathrm{i}\mu\xi$	$\delta(\cdot - \mu)$	1	1	
SαS	$\alpha \in (0,2)$	$-\lvert\xi\rvert^\alpha$	—	α	α	[34]
Sum of SαS	$\alpha, \beta \in (0,2)$	$-\lvert\xi\rvert^\alpha - \lvert\xi\rvert^\beta$	—	$\min(\alpha,\beta)$	$\max(\alpha,\beta)$	[34]
Laplace	—	$-\log(1+\xi^2)$	$\frac{1}{2}\mathrm{e}^{-\lvert x\rvert}$	2	0	[23]
Sym-gamma	$c > 0$	$-c\log(1+\xi^2)$	—	2	0	[23]
Poisson	$\lambda > 0$	$\lambda(\mathrm{e}^{\mathrm{i}\xi} - 1)$	—	2	0	
compound-Poisson	$\lambda > 0, \mathbb{P}_J$	$\lambda(\widehat{\mathbb{P}}_J(\xi) - 1)$	—	variable	0	[47]
Inverse Gaussian	—	—	$\frac{\mathrm{e}^{-x}}{\sqrt{2\pi x^3/2}}\mathbb{1}_{x\geq 0}$	2	1/2	[2]

According to Theorem 3, we have that $\mu_p(w) \leq \frac{d}{\min(p,\beta_0)}$. We conjecture that

$$\mu_p(w) = \frac{d}{\min(p, \beta_0)} \tag{104}$$

for every white noise w.

If this conjecture is true, then $\mu_\infty(w) = 1/\beta_0$. When β_0 goes to 0, we need stronger and stronger weights to include the Lévy white noise into the corresponding Hölder space. The limit case is $\beta_0 = 0$ for which we only have local results. Indeed, polynomial weights are not increasing fast enough to compensate the erratic behavior of the white noise. This is consistent with the fact that a Lévy white noise with $\beta_0 = 0$ is not tempered [9].

8.2 Besov Regularity of Some Specific Lévy White Noises

We apply our results to specific Lévy white noises. We start by recalling the Blumenthal-Getoor indices of the considered white noises. We gives in Table 1 the Lévy exponent and the probability density of the underlying infinitely divisible law, when they can be expressed in a closed form. All the considered distributions are known to be infinitely divisible. For Gaussian, SαS, or compound-Poisson noises, this can be easily seen from the definition. For the others, it is a non-trivial fact, and we refer to [35] for more details and references to the adequate literature.

We moreover remark that any combination of β_0 and β_∞ is possible, as stated in Proposition 10.

Proposition 10. *For every $\beta_0, \beta_\infty \in [0,2]$, there exists a Lévy white noise with Blumenthal-Getoor indices β_0 and β_∞.*



(a) Gaussian (b) Cauchy (SαS, $\alpha = 1$)

(c) Sym-gamma (d) compound-Poisson

(e) Inverse Gaussian

Figure 2: Besov localization of specific Lévy white noises. A white noise is almost surely in a given Besov space $B^{\tau,\mathrm{loc}}_{p,q}(\mathbb{R}^d)$ if $(1/p, \tau)$ is located in the shaded region.

Proof. We first recall that a Lévy exponent can be uniquely represents by its Lévy triplet (μ, σ^2, ν) as [35, Theorem 8.1]

$$f(\xi) = \mathrm{i}\mu\xi - \frac{\sigma^2\xi^2}{2} + \int_{\mathbb{R}\setminus\{0\}} (\mathrm{e}^{\mathrm{i}\xi x} - 1 - \mathrm{i}\xi x \mathbb{1}_{|x|\leq 1})\nu(\mathrm{d}x), \tag{105}$$

where $\mu \in \mathbb{R}$, $\sigma^2 \geq 0$, and ν is a Lévy measure, that is, a measure on $\mathbb{R}\setminus\{0\}$ such that $\int_{\mathbb{R}\setminus\{0\}} \inf(1, x^2)\nu(\mathrm{d}x) < \infty$. When $\mu = \sigma^2 = 0$ and ν is symmetric, the Lévy white noise with Lévy triplet $(0, 0, \nu)$ has Blumenthal-Getoor indices given by [10, Section 3.1]

$$\begin{aligned} \beta_\infty &= \inf_{p\in[0,2]} \left\{ \int_{|x|\leq 1} |x|^p \nu(\mathrm{d}x) < \infty \right\}, \\ \beta_0 &= \sup_{p\in[0,2]} \left\{ \int_{|x|> 1} |x|^p \nu(\mathrm{d}x) < \infty \right\}. \end{aligned} \tag{106}$$

For $0 < \beta_0 \leq 2$ and $0 \leq \beta_\infty < 2$, we set

$$\nu_{\beta_0}(x) = |x|^{-(\beta_0+1)} \mathbb{1}_{|x|>1}, \quad \nu^{\beta_\infty}(x) = |x|^{-(\beta_\infty+1)} \mathbb{1}_{|x|\leq 1}.$$

Moreover, for $\beta_0 = 0$ and $\beta_\infty = 2$, we set

$$\nu_0(x) = \log^{-2}(1 + |x|)|x|^{-1}\mathbb{1}_{|x|>1}, \quad \nu^2(x) = \log^{-1/2}(1 + |x|)|x|^{-3}\mathbb{1}_{|x|\leq 1}.$$

For $0 \leq \beta_0, \beta_\infty \leq 2$ and defining $\nu_{\beta_0}^{\beta_\infty} = \nu_{\beta_0} + \nu^{\beta_\infty}$, we see easily that

$$\int_{\mathbb{R}\setminus\{0\}} \inf(1, x^2)\nu_{\beta_0}^{\beta_\infty}(\mathrm{d}x) < \infty,$$

so that $\nu_{\beta_0}^{\beta_\infty}$ is a Lévy measure. Based on (106), we also see that the associated indices are β_0 and β_∞. □

8.3 Conclusion

We derived new results on the Besov regularity of general multidimensional Lévy white noises. We have shown that the local regularity depends on the Blumenthal-Getoor index β_∞ which measures the behavior of the Lévy exponent at infinity. The global regularity over the complete space $\mathbb{R}^d$ requires the introduction of polynomial weights. We have shown that the second Blumenthal-Getoor index β_0 characterizes the weight parameter. Finally, we applied our results to specific white noises.

Acknowledgment

The authors are grateful to John Paul Ward for the fruitful preliminary discussions that lead to this work. The research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013) / ERC grant agreement n° 267439.

References

[1] D. Applebaum. *Lévy processes and stochastic calculus*. Cambridge university press, 2009.

[2] O.E. Barndorff-Nielsen. Processes of normal inverse Gaussian type. *Finance and Stochastics*, 2(1):41–68, 1997.

[3] Á. Bényi and T. Oh. Modulation spaces, Wiener amalgam spaces, and Brownian motions. *Advances in Mathematics*, 228(5):2943–2981, 2011.

[4] R.M. Blumenthal and R.K. Getoor. Sample functions of stochastic processes with stationary independent increments. *Journal of Mathematics and Mechanics*, 10:493–516, 1961.

[5] V.I. Bogachev. *Measure theory*, volume 1. Springer Science & Business Media, 2007.

[6] B. Böttcher, R. Schilling, and J. Wang. *Lévy Matters III*. Springer, 2013.

[7] P. Cartier. Processus aléatoires généralisés. *Séminaire Bourbaki*, 8:425–434, 1963.

[8] Z. Ciesielski, G. Kerkyacharian, and B. Roynette. Quelques espaces fonctionnels associés à des processus gaussiens. *Studia Mathematica*, 107(2):171–204, 1993.

[9] R.C. Dalang and T. Humeau. Lévy processes and Lévy white noise as tempered distributions. *arXiv preprint arXiv:1509.05274*, 2015.

[10] C.S. Deng and R.L. Schilling. On shift Harnack inequalities for subordinate semigroups and moment estimates for Lévy processes. In *Stochastic Processes and their Applications*, volume 125, pages 3851–3878, 2015.

[11] D.E. Edmunds and H. Triebel. *Function Spaces, Entropy Numbers, Differential Operators*, volume 120. Cambridge University Press, 2008.

[12] J. Fageot, A. Amini, and M. Unser. On the continuity of characteristic functionals and sparse stochastic modeling. *Journal of Fourier Analysis and Applications*, 20:1179–1211, 2014.

[13] J. Fageot, M. Unser, and J.P. Ward. On the Besov regularity of periodic Lévy noises. *Applied and Computational Harmonic Analysis*, in press.

[14] X. Fernique. Processus linéaires, processus généralisés. *Annales de l'Institut Fourier*, 17:1–92, 1967.

[15] I.M. Gelfand. Generalized random processes. *Doklady Akademii Nauk SSSR*, 100:853–856, 1955.

[16] I.M. Gel'fand and N.Y. Vilenkin. *Generalized Functions. Vol. 4: Applications of Harmonic Analysis*. Translated by A. Feinstein. Academic Press, New York-London, 1964.

[17] L. Grafakos. *Classical and Modern Fourier Analysis*. Pearson Education, Inc., Upper Saddle River, NJ, 2004.

[18] V. Herren. Lévy-type processes and Besov spaces. *Potential Analysis*, 7(3):689–704, 1997.

[19] T. Hida and Si-Si. *An Innovation Approach to Random Fields*. World Scientific Publishing Co., Inc., River Edge, NJ, 2004.

[20] R.A. Horn. Quadratic forms in harmonic analysis and the Bochner-Eberlein theorem. *Proceedings of the American Mathematical Society*, 52(1):263–270, 1975.

[21] K. Itô. Stationary random distributions. *Kyoto Journal of Mathematics*, 28(3):209–223, 1954.

[22] K. Itô. *Foundations of Stochastic Differential Equations in Infinite Dimensional Spaces*, volume 47. SIAM, 1984.

[23] S. Koltz, T.J. Kozubowski, and K. Podgorski. *The Laplace Distribution and Generalizations*. Boston, MA: Birkhauser, 2001.

[24] F. Kühn. Existence and estimates of moments for Lévy-type processes. *arXiv preprint arXiv:1507.07907*, 2015.

[25] S. Kusuoka. The support property of a Gaussian white noise and its applications. *Journal of the Faculty of Science. University of Tokyo. Section IA. Mathematics*, 29(2):387–400, 1982.

[26] G. Laue. Remarks on the relation between fractional moments and fractional derivatives of characteristic functions. *Journal of Applied Probability*, pages 456–466, 1980.

[27] H. Luschgy and G. Pagès. Moment estimates for Lévy processes. *Electronic Communications in Probability*, 13:422–434, 2008.

[28] M. Matsui and Z. Pawlas. Fractional absolute moments of heavy tailed distributions. *arXiv preprint arXiv:1301.4804*, 2013.

[29] R.A. Minlos. Generalized random processes and their extension in measure. *Trudy Moskovskogo Matematicheskogo Obshchestva*, 8:497–518, 1959.

[30] D. Nualart and Y. Ouknine. Besov regularity of stochastic integrals with respect to the fractional Brownian motion with parameter h$>1/2$. *Journal of Theoretical Probability*, 16(2):451–470, 2003.

[31] M. Reed and B. Simon. *Methods of Modern Mathematical Physics: Functional Analysis*, volume 1. Gulf Professional Publishing, 1980.

[32] M. Rosenbaum. First order p-variations and Besov spaces. *Statistics & Probability Letters*, 79(1):55–62, 2009.

[33] B. Roynette. Mouvement Brownien et espaces de Besov. *Stochastics: An International Journal of Probability and Stochastic Processes*, 43(3-4):221–260, 1993.

[34] G. Samorodnitsky and M.S. Taqqu. *Stable non-Gaussian processes: Stochastic models with infinite variance*. Chapman and Hall, 1994.

[35] K.-I. Sato. *Lévy processes and infinitely divisible distributions*, volume 68 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2013. Translated from the 1990 Japanese original, Revised edition of the 1999 English translation.

[36] R.L. Schilling. On Feller processes with sample paths in Besov spaces. *Mathematische Annalen*, 309(4):663–675, 1997.

[37] R.L. Schilling. Growth and Hölder conditions for the sample paths of Feller processes. *Probability Theory and Related Fields*, 112(4):565–611, 1998.

[38] R.L. Schilling. Function spaces as path spaces of Feller processes. *Mathematische Nachrichten*, 217(1):147–174, 2000.

[39] H.-J. Schmeisser and H. Triebel. *Topics in Fourier Analysis and Fonction Spaces*. Wiley Chichester, 1987.

[40] L. Schwartz. *Théorie des distributions*. Hermann, 1966.

[41] B. Simon. Distributions and their Hermite expansions. *Journal of Mathematical Physics*, 12(1):140–148, 2003.

[42] F. Trèves. *Topological Vector Spaces, Distributions and Kernels*. Academic Press, New York-London, 1967.

[43] H. Triebel. *Interpolation Theory, Function Spaces, Differential Operators*. North Holland, 1978.

[44] H. Triebel. *Function Spaces and Wavelets on Domains*. Number 7. European Mathematical Society, 2008.

[45] Hans Triebel. *Theory of Function Spaces*. Modern Birkhäuser Classics. Birkhäuser/Springer Basel AG, Basel, 2010. Reprint of 1983 edition [MR0730762].

[46] M. Unser and P.D. Tafti. Stochastic models for sparse and piecewise-smooth signals. *IEEE Transactions on Signal Processing*, 59(3):989–1006, 2011.

[47] M. Unser and P.D. Tafti. *An Introduction to Sparse Stochastic Processes*. Cambridge University Press, Cambridge, UK, 2014.

[48] M. Veraar. Regularity of Gaussian white noise on the d-dimensional torus. *arXiv preprint arXiv:1010.6219*, 2010.